



*Recombinant Human IL-17AF*  
(Interleukin-17AF Heterodimer)

*Catalog Number: 100-161*  
*Accession Number: Q16552/Q96PD4*

*Specifications and Uses:*

**Alternate Names:** None

**Description:**

Interleukin-17AF (IL-17AF) is a member of the IL-17 family of proteins produced by a subset of T cells, called Th17, following stimulation with IL-23. Since IL-17AF is thought to signal through the IL-17RA receptor, its biological function is similar to that of IL-17A in that it induces the production of a variety of chemokines, in addition to airway neutrophilia. In regard to these functions, IL-17AF has less activity than the IL-17A homodimer but, greater activity than the IL-17F homodimer. Human and rat IL-17AF both show activity on mouse cells. Recombinant human IL-17AF is a non-glycosylated heterodimer, containing one IL-17A subunit and one IL-17F subunit. The dimer has a total of 271 amino acids, with an approximate molecular weight of 30.7 kDa.

**Source:** *E.coli*

**Physical Appearance:** Sterile filtered white lyophilized (freeze-dried) powder.

**Formulation and Stability:**

Recombinant human IL-17AF is lyophilized with no additives.

Lyophilized product is very stable at -20°C. Reconstituted material should be aliquoted and frozen at -20°C. It is recommended that a carrier protein (0.1% HSA or BSA) is added for long term storage.

**Reconstitution:**

Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/mL, which can be further diluted into other aqueous solutions.

**Protein Content and Purity (typically ≥ 98%) determined by:**

Reducing and Non-reducing SDS-PAGE, UV spectroscopy at 280 nm

**Endotoxin Level:**

Measured by kinetic LAL analysis and is typically ≤ 1 EU/μg protein.

**Biological Activity:**

The activity is determined by a dose-dependent production of IL-6 in cultured mouse NIH 3T3 fibroblasts and is typically 3-15 ng/mL.

**AA Sequence:**

IL-17A: MIVKAGITIP RNP GCPNSED KNFPRTVMVN LNIHNRNTNT NPKRSSDYYN RSTSPWNLHR  
NEDPERYPSV IWEAKCRHLG CINADGNVDY HMNSVPIQQE ILVLRREPPH CPNSFRLEKI LVSVGCTCVT  
PIVHHVA

IL-17F: MRKIPKVGHT FFQKPESCPP VPGGSMKLDI GIINENQRVS MSRNIESRST SPWNYTVTWD PNRYPSEVVQ  
AQCRNLGCIN AQ

**THIS PRODUCT IS FOR RESEARCH USE ONLY AND IS NOT FOR USE IN HUMANS!**

Gentaur Molecular Products  
Voortstraat 49  
1910 Kampenhout, Belgium