

RayBiotech, Inc.

3607 Parkway Lane suite 100
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

Recombinant Human B-Cell Leukemia / Lymphoma 2 (minus BH3 domain)

Catalog No.
228-10100

Source:
Escherichia coli.

Synonyms:

Apoptosis regulator Bcl-2, BCL2, B-cell CLL/lymphoma 2, Bcl-2.

Introduction:

BCL2 gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends.

Description:

Bcl-2 Des BH3 domain (93-107 residues) Human Recombinant produced in E. coli is a single, non-glycosylated polypeptide chain containing 203 amino acids 1-92 and 108-218.

The Bcl-2 is expressed as His-Tag fusion protein and purified by proprietary chromatographic techniques.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

The protein contains 10mM Tris-HCL pH-8, 1mM EDTA and 250mM NaCl.

Solubility:


Suspend Bcl-2 in 100µl of 0.5M Acetic acid, overnight at 4°C.

Dilute 10 fold into selected buffer system.

BCL-2 has tendency to form intramolecular disulfide bond, 5mM DTT is recommended in assay buffer.

When running SDS-PAGE gel, 10mM DTT is recommended.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 100
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Stability:

Lyophilized Bcl-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Bcl-2 should be stored at 4°C between 2-7 days and for future use below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

Purity:

Greater than 95.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

Applications

Input marker or positive control (Western Blotting).

Function study.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**