

• Rabbit Anti-Integrin alpha 4/CD49d/Integrin α 4 Polyclonal Antibody

Primary Antibodies

Background:

Integrin alpha 4 (also called CD49d) is a 150 kDa protein that possesses a large extracellular domain involved in ligand binding, a single transmembrane domain, and an intracellular regulatory domain possessing multiple sites for phosphorylation. Integrin alpha 4 forms heterodimers with integrins beta 1 and beta 7. Integrin alpha 4 is expressed on leukocytes and leukocyte precursors, neural crest cells, and developing skeletal muscles and is essential for embryogenesis, hematopoiesis, and immune responses. The presence of integrin alpha 4 promotes cell migration and inhibits cell spreading and contractility. Integrin alpha 4 function has been implicated in the pathogenesis of multiple diseases including asthma, rheumatoid arthritis, Crohn's disease, ulcerative colitis, hepatitis C, and multiple sclerosis, and therefore, modulation of integrin alpha 4 function has become an important target for drug discovery.

Source/Purification:

KLH conjugated synthetic peptide derived from human Integrin alpha 4 C-terminus. Was purified by Protein A and peptide affinity chromatography.

Storage: Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Cow, Horse, Rabbit,

Application:

- WB(1:100-500)
- ELISA(1:500-1000)
- IP(1:20-100)
- IHC-P(1:100-500)
- IHC-F(1:100-500)
- FACS(1:100-500)
- IF(1:100-500)
- Not yet tested in other applications.
Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 114kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.