

bs-0784R-PE-Cy3

• Rabbit Anti-IFN- β Polyclonal Antibody, PE-Cy3 conjugated

Conjugated Primary Antibodies

Background:

The mammalian type I Interferons (IFN1) are produced in response to viral infection and other inducers. They are divided into alpha and beta subtypes leukocytes and fibroblasts reactivity. The human IFN alphas are encoded by a family of at least 15 different genes, while IFN beta is the unique member of its subtype. There is approximately 50% amino acid homology between the alpha and beta subtypes. Both IFN subtypes are pleiotropic cytokines and have a similar range of biological activities. Differences between alpha subtypes, and between IFN alpha and betas, are in potency and cell type specific activities. In particular, IFN beta elicits a markedly higher antiproliferation response in some cell types such as, embryonal carcinoma, melanoma and melanocytes than do IFN alphas. Higher potency of IFN beta in treatment of multiple sclerosis and certain cancers has been observed. Type I IFNs signal through binding to a common cell surface receptor. Two chains of the receptor, IFNAR1 and IFNAR2, have been identified. Both chains are necessary for function and in the absence of either there is neither high affinity binding nor biological activity. The intracellular portions of the receptor subunits are bound by tyrosine kinases, Jak1 and Tyk2, members of the Janus kinase family. Upon ligand binding these kinases are activated and phosphorylate members of the STAT family of transcription factors, as well as IFNAR1 and 2.

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities: Mouse,Rat,

Application:

- IF(1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 20kDa

Preservatives:

10ug/uL BSA and 0.1% NaN₃.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.

For full size images and description please click [HERE](#).