

bs-1173R-A555

• Rabbit Anti- TNFRSF18/GITR Polyclonal Antibody, Alexa Fluor 555 conjugated

Conjugated Primary Antibodies

Background:

TNFRSF18 or GITR (glucocorticoid-induced TNF receptor family-regulated gene) is a 25 kD TNF receptor superfamily member (also known as AITR and TNFRSF18) originally identified in the mouse by comparing untreated and dexamethasone-treated murine T cell hybridoma cells. Human GITR was subsequently identified by searching an expressed sequence tag database. GITR is expressed on activated lymphocytes and is upregulated by T cell receptor engagement. The cytoplasmic domain of GITR is homologous to CD40, 4-1BB and CD27 and has been shown to interact with TRAF 1-3, but not TRAF 5 or 6. GITR signaling has been shown to regulate T cell proliferation and TCR-mediated apoptosis, and to break immunological self-tolerance. GITR appears to be highly expressed on CD4+CD25+ T regulatory cells and has been shown to induce NF- κ B activation through TRAF2/NIK signaling. GITR has been proposed to be involved in the development of regulatory T cells and to regulate the activity of Th1 subsets. GITR binds GITRL, a TNF superfamily ligand expressed on endothelial cells, dendritic cells, macrophages, and B cells.

Purification: Was purified by Protein A and peptide affinity chromatography.



Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.



For full size images and description please click [HERE](#).

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities: Human, Mouse, Rat,

Application:

- IF(1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 26kDa

Preservatives:

10ug/uL BSA and 0.1% NaN₃.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.