

bs-6353R-PE-Cy3

• Rabbit Anti-HIPK2 Polyclonal Antibody, PE-Cy3 conjugated

Conjugated Primary Antibodies

Background:

Protein kinase acting as a corepressor of several transcription factors, including SMAD1 and POU4F1/Brn3a and probably NK homeodomain transcription factors. Inhibits cell growth and promotes apoptosis. Involved in transcriptional activation of TP53 and TP73. Phosphorylation of TP53 may be mediated by a TP53-HIPK2-AXIN1 complex. In response to TGFB, cooperates with DAXX to activate JNK. Phosphorylates the antiapoptotic factor CTBP1 and promotes its proteasomal degradation. In the Wnt/beta-catenin signaling pathway acts as an intermediate kinase between TAK1 and NLK to promote the proteasomal degradation of MYB (By similarity). Phosphorylates CBX4 upon DNA damage and promotes its E3 SUMO-protein ligase activity. PML, HIPK2 and FBXO3 may act synergically to activate p53/TP53-dependent transactivation.

Purification: Was purified by Protein A and peptide affinity chromatography.


Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.



For full size images and description please click [HERE](#).

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Chicken, Dog, Pig, Rabbit,

Application:

- IF(1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 132kDa

Preservatives:

10ug/uL BSA and 0.1% NaN₃.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.