

bs-7111R-PE

• Rabbit Anti-EDA2R/XEDAR Polyclonal Antibody, PE conjugated

Conjugated Primary Antibodies

Background:

The tumor necrosis factor receptor (TNFR) superfamily represents a growing family of type I transmembrane glycoproteins that are involved in various cellular functions, including proliferation, differentiation and programmed cell death (1–3). These proteins share homology for cysteine-rich repeats in the extracellular ligand binding domain and an intracellular death domain (1–3). Members of the TNFR superfamily transmit signals through protein-protein interactions, and these signals can lead to the activation of either the caspase and Jun kinase pathways, which promote cell death, or the NFκB pathway, which results in cell survival (1). The ectodermal dysplasia receptor (EDAR) promotes all three of these pathways and mediates ectodermal differentiation (4). EDAR is encoded by the downless gene and is mutated in ectodermal dysplasia syndromes, which are characterized by impaired hair, teeth and sweat gland development (5). Ectodysplasin A (EDA) is a type II membrane protein that is encoded by the Tabby gene and produces many splice variants, the longest of which, EDA-A1, serves as the ligand for EDAR (5–7). EDA-A2, which differs from EDA-A1 by the deletion of two amino acids, binds only the X-linked ectodysplasin-A2 receptor (XEDAR) (7). Both EDAR and XEDAR exhibit homology with TROY (8).

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities: Human, Mouse, Rat,

Application:

- IF (1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 33kDa

Preservatives:

10ug/uL BSA and 0.1% NaN₃.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.

For full size images and description please click [HERE](#).